



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

40. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Find the law of the force, in order that the orbit may be a Cassinian Oval.

Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

Let $r^4 + 2c^2r^2\cos 2\theta = m^4 - c^4 = a^4 \dots (1)$ be the equation to the oval.

Then $a^4u^4 = 2c^4u^2\cos 2\theta + 1$, where $u = 1/r$.

$$\frac{du}{d\theta} = \frac{c^2u\sin 2\theta}{c^2\cos 2\theta - a^4u^2} \dots (2).$$

$$\frac{d^2u}{d\theta^2} = \frac{\left\{ (c^2\sin 2\theta \frac{du}{d\theta} + 2c^2u\cos 2\theta)(c^2\cos 2\theta - a^4u^2) + (2c^2\sin 2\theta + 2a^4u \frac{du}{d\theta})c^2u\sin 2\theta \right\}}{(c^2\cos 2\theta - a^4u^2)^2}$$

$$= \left\{ \frac{3c^4r^8 + 10r^4c^4m^4 - 3r^4c^8 - (m^4 - c^4)^3 - 7r^4m^8 + 9r^8m^4 - r^{12}}{r(a^4 + 2r^4)^3} \right\}.$$

$$F = \text{force} = h^2u^2(u + \frac{d^2u}{d\theta^2}) = \frac{h^2}{r^3} + \frac{h^2}{r^2} \cdot \frac{d^2u}{d\theta^2}.$$

$$\therefore F = \frac{h^2(7r^{12} + 21r^8m^4 + 3r^4c^8 - 9c^4r^8 - 2c^4m^4r^4 - m^8r^4)}{r^3(m^4 - c^4 + 2r^4)^3}$$

$$= \frac{h^2\{7m^9 + 21m^5r^5 - 9c^4r^5 + 3rc^8 - 2c^4m^4r - m^8r\}}{(m^4 - c^4 + 2r^4)^3}.$$

41. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics, Columbian University, Washington, D. C.

If the earth were a perfect sphere and had a frictionless surface, what would be the motion of a ball placed at a given latitude?

[No solution of this problem has been received. EDITOR].

42. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics, Columbian University, Washington, D. C.

Find the time of vibration of a particle slightly displaced from the center of a solid cylinder in direction of the axis, the matter of the cylinder attracting according to the laws of nature.